

## Bionomics of Polyphagous Red Spider Mite, *Tetranychus urticae* Koch Under *In-vitro* Conditions

S. Shanchitha\*, N. Thiruchchelvan and G. Mikunthan

Department of Agricultural Biology, Faculty of Agriculture, University of Jaffna, Sri Lanka

\*shanchitha98@gmail.com

The red spider mite (*Tetranychus urticae* Koch) is a substantial agricultural pest with a broad host range and high reproductive potential. This study explores the biological traits such as development stages, reproductive biology and morphology of red spider mite under laboratory conditions ( $31 \pm 1^\circ\text{C}$  and  $54 \pm 3\%$  RH). The red spider mite was reared on a 2.5 cm leaf disc array set up in a petri dish. Reading was taken at four-hour intervals to measure the development period. The results revealed that, its life cycle comprised egg, larva, protonymph, deutonymph, and adult stages. The immature stages were distinguished by brief periods of quiescence known as nymphochrysalis, dutochrysalis, and teliochrysalis. The male and female had recorded incubation periods of  $65.07 \pm 1.83$  and  $64 \pm 0$  hours, larval periods of  $13.60 \pm 2.53$  and  $16.80 \pm 3.10$  hours, protonymphal periods of  $8.53 \pm 1.41$  and  $11.47 \pm 2.10$  hours, and deutonymphal periods of  $12 \pm 0$  and  $14.93 \pm 2.37$  hours, respectively. Male development took shorter ( $140.27 \pm 4.40$  hours) than female ( $151.20 \pm 3.10$  hours) from egg to adult emergence. On average,  $122.10 \pm 19.77$  and  $85 \pm 18.56$  eggs were produced by mated and unmated females, respectively. Sexual dimorphism was observed in the deutonymphal stage, as the body size of females ( $0.34 \pm 0.02$  mm length and  $0.27 \pm 0.02$  mm width) was higher than that of males ( $0.30 \pm 0.02$  mm length and  $0.20 \pm 0.02$  mm width). Studying the bionomics of red spider mites under in-vitro conditions would be helpful to set the optimum conditions for rearing or mass multiplication of red spider mites for various research studies.

**Keywords:** Development period, Life cycle, Morphology, Reproductive biology, *Tetranychus urticae*